

TURI SURFACE SOLUTIONS LABORATORY EVALUATION SUMMARY

DateRun: 10/22/2014
Experimenters: Liang; Nguyen; Curtis; Langberg;
ClientType: Pollet Odor Elimination
ProjectNumber: 1
Substrates: Glass Bottle;
Contaminants: Milk;
CleaningMethods: Spray;
AnalyticalMethods: Odor panel;
Purpose: To evaluate supplied products for odor elimination
ExperimentalProcedure: 250ml glass bottles were filled with 1ml of milk each and left to spoil over a 3 day period. A panel of three then examined the odors to determine baseline values using 6 bottles, including a control. The bottles were then treated with the cleaners at the recommended dilutions. Each panelist was asked to describe odor and rank the level of intensity of the malodor 1 being the worst smell and 5 being the best. After the panelists observed the initial odors, bottles were recapped and observations were recorded. Bottles were reopened and more cleaners were applied. Each bottle was subjected to additional rounds of treatment and each panelist was used to assess malodor levels. Additional observations were made the following day to check malodor elimination efficiency. When the intensity of malodor level has reached a rating of 4 or higher; any additional sprays in the contaminated bottles are unnecessary as the malodor level is no longer noticeable. On the 4th day of the test; an additional test was conducted to see if the contaminated bottles have reached back to its original malodor level. Only contaminated bottles that indicated an increase in malodor level of a rating below a 4 was retested with two more sprays to see if the cleaner can effectively remove the increased overnight malodor level.

ChemistriesEvaluated: Polbio Enzytan; Febreze;

Results: Each of the three panelists observed decreases in the malodor. The non-treated sample was nearly unchanged from the start of the testing.

Cleaner	Tester #	# Of Sprays	Bottle 1	Bottle 2	Bottle 3	Average
Polbio Enzytan	1	0	1	1	1	1.0
		2	3.5	3.2	4	3.6
		4	3.8	4.5	4	4.1
		6	N/A	N/A	N/A	
		8	N/A	N/A	N/A	
	2	0	1	1	1	1.0
		2	5	4	4	4.3
		4	N/A	N/A	N/A	
		6	N/A	N/A	N/A	
		8	N/A	N/A	N/A	
	3	0	1	1	1	1.0
		2	3.5	3	4	3.5
		4	4.5	4	4	4.2
		6	N/A	N/A	N/A	
		8	N/A	N/A	N/A	
Febreze	1	0	1	1	1	1.0
		2	2.6	3	2.9	2.8
		4	3	3.5	3.5	3.3
		6	3.2	3.5	3.5	3.4
		8	N/A	N/A	N/A	
	2	0	1	1	1	1.0
		2	1.5	3	2	2.2
		4	2.5	3	2.5	2.7
		6	4	4	3.5	3.8
		8	N/A	N/A	N/A	
	3	0	1	1	1	1.0

	2	2.5	2	2.5	2.3
	4	3	3	3	3.0
	6	3	3	3	3.0
	8	N/A	N/A	N/A	

4th Day

Cleaner	Tester #	# Of Sprays	Bottle 1	Bottle 2	Bottle 3	Average
Polbio Enzysan	1	0	3.2	3.6	3.4	3.4
		2	3.8	4	4	3.9
	2	0	3.5	3.5	3	3.3
		2	3.8	4.1	3.9	3.9
	3	0	3.8	3.5	2.5	3.3
		2	3.7	4	4.1	3.9
Febreeze	1	0	2	2	2.7	2.2
		2	2.5	2.8	2.7	2.7
	2	0	2	2.5	2.5	2.3
		2	2.5	2	2.3	2.3
	3	0	2	2	2	2.0
		2	2.6	2	2.8	2.5

Summary

Substrates: Glass
Contaminants: Milk

Summary	Control	Polbio Enzysan	Febreeze
Original (day 3)	1	1	1
2 Spray	1	3.8	2.4
4 Spray	1	4.1	3
6 Spray	1	N/A	3.4
8 Spray	1	N/A	N/A

Conclusion:

Based on our observed results Polbio Enzysan was better at odor elimination than Febreeze. Polbio was able to almost completely eliminate the milk malodor with the initial 2 sprays. Polbio eliminated milk malodor within 4 sprays while Febreeze was not able to even after 8 sprays. On the 4th day observations showed that Polbio still was effective at malodor elimination but Febreeze had declined in effectiveness.